



Gyrokinetic linearized Landau collision operator

Madsen, Jens

Publication date:
2014

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Madsen, J. (2014). *Gyrokinetic linearized Landau collision operator*. Poster session presented at 15th European Fusion Theory Conference, Oxford, United Kingdom.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

ABSTRACT

Here the full gyrokinetic electrostatic linearized Landau collision operator, entering the δF formulation of gyrokinetic theory, is calculated including the equilibrium operator $\mathcal{C}[F_{Ma}, F_{Mb}]$. Energy exchange between plasma species is described by $\mathcal{C}[F_{Ma}, F_{Mb}]$.

Furthermore, $\mathcal{C}[F_{Ma}, F_{Mb}]$ describes drag and diffusion of the magnetic field aligned component of the vorticity associated with the $\mathbf{E} \times \mathbf{B}$ -drift.

Contrary to conventional wisdom it is shown that $\mathcal{C}[F_{Ma}, F_{Ma}] \neq 0$ for **like-particle collisions** and has the same order of magnitude as the gyrokinetic test- and field-particle operators. $\mathcal{C}[F_{Ma}, F_{Ma}]$ **must be included in gyrokinetic δF collision operators**.

PARTICLES COLLIDE

Gyrokinetic theory decouples the fast time-scale associated with the fast cyclotron motion of charged particles in strongly magnetized plasmas from the low-frequency dynamics of gyro-centers. The gyrokinetic theory describes the time-evolution of gyro-centers.

A gyro-center is a mathematical construction. Gyro-centers do not carry charge nor do they collide. Therefore, caution must be exercised when describing particle dynamics in the gyro-center picture.

For instance in the gyrokinetic Gauss's Law the charge distribution must be deduced from the distribution of gyro-centers. Therefore, the gyro-center charge contribution is accompanied by polarization density and inverse gyro-angle averages, which in combination give the low-frequency particle charge distribution. **The polarization density originates from the Maxwellian (constant density) part of the gyrokinetic distribution function.**

Similarly, the Landau collision operator (and collisions in general) describes collisions between particles, not gyro-centers. Therefore, collision operators in gyrokinetic theory are expected to have terms similar to those in the gyrokinetic Gauss's Law. **A polarization-density-like contribution is expected in gyrokinetic collision operators.**

GYROKINETIC MAXWELLIAN

The Landau collision operator vanishes $\mathcal{C}[f_a, f_b] = 0$ when $f_{a,b}(\mathbf{x}, \mathbf{v})$ are Maxwellian $f_M = n_0(2\pi m T_0)^{-3/2} \exp(-\frac{m\mathbf{v}^2}{2T_0})$, with the same temperature. However, when evaluating the collision operator with two gyrokinetic Maxwellians: $F_{Ma} = N(2\pi T m_a)^{-3/2} \exp(-\frac{1/2 m_a v_{\parallel}^2 + \mu B}{T})$ having the same temperature the Landau collision operator **does not vanish**:

$$\mathcal{C}[F_{Ma}, F_{Mb}] \neq 0.$$

f_M and F_M have the same functional form but do only agree to lowest order in ϵ :

$$F_M(\mathbf{Z}) = f_M(\mathbf{z})[1 - \epsilon q \tilde{\phi}(\mathbf{z})/T] + \mathcal{O}(\epsilon^2).$$

REFERENCES

- [1] M. N. Rosenbluth et al, *Phys. Rev.*, 1, 107 (1957).
- [2] Jens Madsen. *Phys. Rev. E*, 87, 011101 (2013).
- [3] P. Helander and D.J. Sigmar. *Collisional Transport in Magnetized Plasmas*. 2005.
- [4] X. Q. Xu, M. N. Rosenbluth. *Phys. Fluids B*, 3, 627 (1991).
- [5] B. Li and D. R. Ernst. *Phys. Rev. Lett.*, 106, 195002 (2011).

ACKNOWLEDGEMENTS

This work was funded by an EFDA fusion researcher fellowship (WP11-FRF-RISOE/MADSEN)

LANDAU COLLISION OPERATOR

In this work we express the Landau collision operator in gyro-center coordinates and perform an explicit gyro-angle average. In (\mathbf{x}, \mathbf{p}) coordinates the Landau collision operator is

$$\mathcal{C}[f_a; f_b] = \frac{\partial}{\partial p_i} \left[\frac{m_a}{m_b} f_a \frac{\partial H_b}{\partial p_i} - m_a^2 \frac{\partial^2 G_b}{\partial p_i \partial p_j} \frac{\partial f_a}{\partial p_j} \right],$$

where the Rosenbluth potentials[1] are defined as:

$$\begin{pmatrix} H_b \\ G_b \end{pmatrix} = \frac{-\Gamma_{ab}}{8\pi} \int d^6 z' f_b(\mathbf{z}') \delta^3(\mathbf{x} - \mathbf{x}') \begin{pmatrix} 2u^{-1} \\ u \end{pmatrix}.$$

$u = |\mathbf{v} - \mathbf{v}'|$ is the relative velocity and $\Gamma_{ab} = (q_a q_b / \epsilon_0)^2 \ln \Lambda$.

The δF formulation arise when the gyro-angle averaged distribution function is split into an equilibrium part F_M and a perturbed part δF . The bi-linearity of the collision operator leads to:

$$\sum_{a,b} \mathcal{C}[\langle F_a \rangle; \langle F_b \rangle] = \sum_{a,b} \underbrace{\mathcal{C}[F_{Ma}; F_{Mb}]}_{\text{equilibrium-operator}} + \underbrace{\mathcal{C}[\delta F_a; F_{Mb}]}_{\text{test-particle-operator}} + \underbrace{\mathcal{C}[F_{Ma}; \delta F_b]}_{\text{field-particle-operator}}$$

- *Main focus here*[2] is to calculate the gyro-angle average of $\mathcal{C}[F_{Ma}; F_{Mb}]$.
- The equilibrium operator is composed of *test-particle-like* and *field-particle-like* parts:

$$\mathcal{C}[F_{Ma}; F_{Mb}] = \mathcal{C}_M^T + \mathcal{C}_M^F$$

TEST-PARTICLE-LIKE OPERATOR

The gyro-angle average of \mathcal{C}_M^T becomes:

$$\begin{aligned} \langle \mathcal{C}_M^T \rangle = & \int d^3 k e^{i\mathbf{k} \cdot \mathbf{X}} \left\{ \left(1 - \frac{m_a}{m_b} \right) \nu_s F_{aM} \left[\frac{m_a v_0^2}{T_a} + \frac{q_a \rho_0^2 k_{\perp}^2 \phi_k}{2T_a} [J_0(k_{\perp} \rho_0) + J_2(k_{\perp} \rho_0)] \right] + \Gamma_{ab} \frac{m_a}{m_b} F_{Ma} F_{Mb} \right. \\ & + \nu_D F_{aM} \left[-\frac{m_a v_0^2}{T_a} - \frac{q_a \rho_0^2 k_{\perp}^2 \phi_k}{2T_a} \left\{ J_0(k_{\perp} \rho_0) [1 + \frac{2v_0^2}{c_{\perp}^2}] + \frac{J_2(k_{\perp} \rho_0)}{2} [1 - \frac{m_a v_0^2}{T_a}] \right\} \right] \\ & \left. + \nu_{\parallel} F_{aM} \left[\frac{m_a v_0^2}{2T_a} \left[\frac{m_a v_0^2}{T_a} - 1 \right] - \frac{q_a \rho_0^2 k_{\perp}^2 \phi_k}{2T_a} (J_0(k_{\perp} \rho_0) [1 - \frac{m v_0^2}{T_a}] - \frac{J_2(k_{\perp} \rho_0)}{2} [1 + \frac{2m_a v_0^2}{T_a}]) \right] \right\} \end{aligned}$$

where $v_0^2 = v_{\parallel}^2 + 2\mu B/m_a$, and $\nu_s = (m_a^2 v_0)^{-1} dH_{0b}/dv_0$, $\nu_D = -2(m_a^2 v_0^3)^{-1} dG_{0b}/dv_0$, $\nu_{\parallel} = -2(m_a^2 v_0^2)^{-1} d^2 G_{0b}/dv_0^2$ denote the standard slowing-down, deflection and parallel velocity diffusion frequencies[3], respectively.

- The zeroth order terms (independent of ϕ) mainly describe energy exchange.

FIELD-LIKE-OPERATOR

The field-particle like operator is characterized by having the time-dependent electric potential residing in the Rosenbluth potentials. The gyro-angle average of the field-particle-like becomes:

$$\begin{aligned} \langle \mathcal{C}_M^F \rangle = & \int d^3 k e^{i\mathbf{k} \cdot \mathbf{X}} \sum_{n=-\infty}^{\infty} F_{Ma} J_{-n}(k_{\perp} \rho_0) \left[\left(1 - \frac{m_a}{m_b} \right) (2\mu \frac{\partial}{\partial \mu} + v_{\parallel} \frac{\partial}{\partial v_{\parallel}}) \frac{-q_b \phi_k h_{nk}}{m_a T_a} \right. \\ & \left. + \left(\frac{n^2 T_a}{4\mu B} + \frac{2\mu B - T_a}{2B} \frac{\partial}{\partial \mu} + \frac{\mu B (2\mu B - T_a)}{B^2} \frac{\partial^2}{\partial \mu^2} + \frac{2\mu B v_{\parallel} T_a}{B} \frac{\partial^2}{\partial \mu \partial v_{\parallel}} + \frac{m_a v_{\parallel}^2 - T_a}{2m_a} \frac{\partial^2}{\partial v_{\parallel}^2} \right) \frac{2q_b \phi_k g_{nk}}{T_a^2} \right], \end{aligned}$$

where

$$\begin{aligned} \begin{pmatrix} h_{nk} \\ g_{nk} \end{pmatrix} = & -\frac{\Gamma_{ab}}{8\pi} \int d\mu' dv'_{\parallel} 2\pi F_{Mb} \left[[\delta_{n0} - J_0(k_{\perp} \rho'_0) J_n(-k_{\perp} \rho'_0)] \frac{\partial}{\partial \mu'} \right. \\ & \left. - \sum_{l=1}^{\infty} \frac{n}{2l} k_{\perp} \frac{\partial \rho'_0}{\partial \mu'} [J_{n+l}(-k_{\perp} \rho'_0) - J_{n-l}(-k_{\perp} \rho'_0)] \right] \begin{pmatrix} 2u_n^{-1} \\ u_n \end{pmatrix}. \end{aligned}$$

Fortunately h_{nk} and g_{nk} are *time-independent*, and can be precomputed and re-used for all time-steps in numerical simulations.

- The ϕ -dependent terms in \mathcal{C}_M^T and \mathcal{C}_M^F describe viscous damping of the magnetic field aligned component of the $\mathbf{E} \times \mathbf{B}$ -vorticity: $\hat{\mathbf{b}} \cdot \nabla \times \mathbf{u}_E \simeq -k_{\perp}^2 \phi_k / B$.
- In the gyrokinetic ordering \mathcal{C}_M^T and \mathcal{C}_M^F has the same order of magnitude as the standard test- and field-particle operators and must therefore be retained. Similar conclusions hold for model operators.

LIKE-PARTICLE COLLISIONS

Utilizing $\mathcal{C}[f_{Ma}, f_{Ma}] = 0$ allows us to evaluate the equilibrium operator:

$$\mathcal{C}[F_{Ma}, F_{Ma}] = \mathcal{C}[F_{Ma} \frac{q \langle \phi \rangle}{T}, F_{Ma}] + \mathcal{C}[F_{Ma}, F_{Ma} \frac{q \langle \phi \rangle}{T}]$$

using the gyrokinetic test- and field-particle operators[4, 5, 2]. $\mathcal{C}[F_{Ma}, F_{Ma}]$ has same order of magnitude as test- and field-particle operators. **Must be retained in order to describe ion-ion collisions correct.**